

Study of the Drell-Yan process with CMS.


QFTHEP 2013: XXIth International Workshop High Energy Physics and Quantum Field Theory , 23-30 Jun 2013, Saint Petersburg, Saint Petersburg (Russian Federation)

I. Gorbunov, On behalf of the CMS Collaboration

JINR, Dubna

June 24, 2013

CMS Detector



Global muon momentum resolution:

$dp_T/p_T = 1 - 1.5\%$ at $p_T = 10 \text{ GeV}$, $dp_T/p_T = 6 - 17\%$ at $p_T = 1 \text{ TeV}$

Electron energy resolution:

$dE/E < 0.5\%$ for $E > 100 \text{ GeV}$

Measurement of the Drell-Yan Cross Section in pp Collisions at $\sqrt{s} = 7 \text{ TeV}$

Based On: CMS-PAS-SMP-13-003

Available on the CERN CDS information server

CMS PAS SMP-13-003

CMS Physics Analysis Summary

Contact: cms-pag-conveners-smp@cern.ch

2013/04/23

Measurement of the differential and double-differential Drell-Yan cross sections in proton-proton collisions at $\sqrt{s} = 7 \text{ TeV}$


The CMS Collaboration

Abstract

Measurements are presented for the measurements of the differential and double differential Drell-Yan cross section using an integrated luminosity of 4.5 fb^{-1} (dimuon) and 4.8 fb^{-1} (dielectron) of proton-proton collision data recorded with the CMS detector at the LHC at $\sqrt{s} = 7 \text{ TeV}$. The cross sections $d\sigma/dM$ are measured in the mass range 15 GeV to 1500 GeV for the dimuon, dielectron, and combined channels. Results are also presented on the measurement of the double-differential cross section $d^2\sigma/dM dY$ in the dimuon channel. The double-differential cross section measure-

The Drell-Yan process

- The production of lepton pairs in pp-collisions is described by the s-channel exchange of γ^*Z
- Theoretical calculations of $d\sigma/dM(\ell\ell)$ and $d^2\sigma/dM(\ell\ell)Y$ ($M(\ell\ell)$ - dilepton invariant mass, Y - dilepton rapidity) established up to NNLO order
- Comparison of Data and MC provide stringent tests of QCD and significant constraints on the evaluation PDFs
- DY is a major background for $t\bar{t}$ and diboson measurements as well as for searches for new physics (high mass dilepton resonances)



Measurements are done at $\sqrt{s} = 7 \text{ TeV}$ on CMS data corresponding to $L = 4.5 \text{ fb}^{-1}$ (dimuon) and $L = 4.8 \text{ fb}^{-1}$ (dielectron) in $20 < M(\ell\ell) < 1500 \text{ GeV}$ range, results are normalized to the cross section in the Z region ($60 < M(\ell\ell) < 120 \text{ GeV}$)

Cross sections are calculated as



$$\sigma = \frac{N_u}{A\epsilon\rho L'} \quad (1)$$

where N_u is the unfolded background-subtracted yield and A - acceptance, ϵ - efficiency (estimated from simulation). ρ - factor that accounts for differences in the efficiency between data and simulation. L is not required for the measurements since the cross sections are normalized to the Z region.

- Dimuon trigger with minimum p_T threshold
- One muon with $|\eta| < 2.4$ and $p_T > 14 \text{ GeV}$ and one with $p_T > 9 \text{ GeV}$
- Standard CMS muon identification and quality criteria (number of hits etc.)
- Impact parameter in the transverse plane less than 2mm
- Angle between the two muons must differ from π by more than 5 mrad
- Dimuon vertex χ^2 probability smaller than 2%.
- Isolation of both muons
 $I_{rel} = (\sum p_T(\text{tracks}) + \sum E_T(\text{had}))/p_T(\mu) < 0.15$, where
 $\sum E_T(\text{had})$ - sum of the E_T of hadronic deposits and
 $\sum p_T(\text{tracks})$ sum of the p_T of the additional tracker tracks
in cone $\Delta R = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2} < 0.3$



- Dielectron trigger with E_T threshold 17 GeV for one of the electrons and 8 GeV for the other
- Candidate is required to be consistent with a particle originating from the primary vertex
- Electron identification criteria based on shower shape and track-cluster matching are applied
- No missing hits in the pixel detector
- Eliminating those electrons for which a partner track consistent with a conversion hypothesis is found
- Isolation
($\sum p_T(\text{tracks}) + \sum E_T(\text{had}) + \sum E_T(\text{em})$)/ $p_T(e) < 0.15$,
where $\sum E_T(\text{em})$ is the sum of the E_T of electromagnetic deposits, except for electrons with $E_T < 20$ GeV where requirement is less than 0.1
- $|\eta| < 1.44$ or $1.57 < |\eta| < 2.5$
- Electron $E_T > 10$ GeV

Acceptance and Efficiency









DY acceptance (blue, filled circles), efficiency (red, open triangles), and their product (black, open squares) per invariant mass bin, for the $\mu^+\mu^-$ (left) and e^+e^- (right) channels

Background 1




The observed dimuon (left) and dielectron (right) invariant mass spectra within the detector acceptance for data and Monte Carlo events and corresponding data to MC ratio of yields. The FEWZ-POWHEG correction is applied to the Monte Carlo signal events. Electron energy correction is applied to data and simulated events. QCD and $t\bar{t}$ background yields as predicted by data-driven methods.

Background 2









The observed dimuon rapidity spectra per invariant mass slice for data and Monte Carlo events.

Normalized rapidity-invariant mass spectrum





The Drell-Yan rapidity-invariant mass spectrum in detector acceptance, normalized to the Z resonance region, $r = 1/\sigma_Z * d^2\sigma/dM dY$.

Comparison with theory expectations




Comparison with theory expectations with various NNLO PDF sets: CT10, HERA, NNPDF2.1, MSTW08, CT10W, JR09, ABKM.

Normalized invariant mass spectrum



The Drell-Yan invariant mass spectrum in the dimuon (left) and dielectron (right) channel, normalized to the Z region, $r = 1/\sigma_Z * d\sigma/dM$, as measured and as predicted by NNLO, for the full phase space.

Combined DY invariant mass




The Drell-Yan invariant mass spectrum in the combined channel, normalized to the Z region, $r = 1/\sigma_Z * d\sigma/dM$ (NNLO).

Vector and axial-vector currents in the process $q\bar{q} \rightarrow Z/\gamma^* \rightarrow l^+l^-$ lead to P-symmetry breaking and forward-backward asymmetry (A_{FB}) in the number of Drell-Yan lepton pairs:

$$A_{FB} = \frac{\sigma_F - \sigma_B}{\sigma_F + \sigma_B} \quad (2)$$



$$\sigma_F = \int_0^1 \frac{d\sigma}{d(\cos\theta)} d(\cos\theta) \quad (3)$$

$$\sigma_B = \int_{-1}^0 \frac{d\sigma}{d(\cos\theta)} d(\cos\theta) \quad (4)$$



Where θ - is the angle between μ^- and quark momentum in dimuon center of mass frame.

Application of A_{FB}




In case of BSM physics A_{FB} can help to determine it's nature

Measurement of the A_{FB} can provide:

- Test of the SM in the new energy region
- Provide at the Z-pole a precise measurement of $\sin^2 \theta_W$

Deviations from the SM predictions for A_{FB} may indicate the existence of:

- A new neutral gauge boson
- Quark-lepton compositeness
- Existence of supersymmetric particles
- Extra dimensions, etc.



- On pp colliders quark and antiquark directions are unknown
- Dimuon boost direction is assumed to be the quark direction

The Collins-Soper frame where the angle θ^* is defined to be the angle between the lepton momentum and a z-axis that bisects the angle between q and \bar{q} directions is used. This frame reduces the uncertainties due to the transverse momentum of quarks.

$$\frac{d\sigma}{d\cos\theta^*} = A(1 + \cos^2\theta^*) + B\cos\theta^* \quad (5)$$

Main sources of background:

- For low masses
 - $Z \rightarrow \tau\tau$
 - QCD dijets
- $t\bar{t}$ for high masses

A_{FB} measurement lead to a large systematic uncertainty and thus requires correction on the following effects:

- Finite detector resolution and alignment
- QED FSR
- Acceptance
- Unknown quark antiquark direction at the LHC

Selection Cuts for Muon Candidates

Kinematics:

- Oppositely charged global and tracker muons
- $|\eta| < 2.4$ and $p_T > 20 \text{ GeV}$

Quality requirements:

- > 10 tracker hits, > 1 pixel hits, > 1 muon hits
- > 2 muon stations
- $\chi^2 < 10$ for the global fit
- Small impact parameter $d_0 < 0.2$
- Remove back-to-back dimuons $\alpha > 2.5 \text{ mrad}$

Isolation:

- Tracker+HCAL isolation
 $(\sum p_T(\text{tracks}) + \sum E_T(\text{had}))/p_T(\mu) < 0.15$

Selection Cuts for Electron Candidates

Kinematics:

- Oppositely charged electrons
- $|\eta| < 1.444$ or $1.566 < |\eta| < 2.5$
- HLT ECAL L1-trigger requiring an ECAL cluster with minimum E_T between 10 to 17 GeV





Quality requirements:

- Minimum supercluster E_T of 20 GeV
- No missing Tracker hits before the first one
- Reject when a conversion partner track close to the electron
- Using shower shape variables

Isolation:

- Isolation using the Tracker and calorimeters

CMS Results with 5 fb^{-1}



arXiv:1207.3973 ; CMS-EWK-11-004 ; CERN-PH-EP-2012-187. - 2012. - 29 p.

$$\frac{d\sigma_{pp \rightarrow l+l-\chi}(Y, s, \cos \theta^*)}{dY ds d\cos \theta^*} \propto \sum_{q=u,d,s,c,b} [\hat{\sigma}_{qq}^{even}(s, \cos^2 \theta^*, \sin^2 \theta_W) + D_{q\bar{q}}(s, Y) \times \hat{\sigma}_{q\bar{q}}^{odd}(s, \cos \theta^*, \sin^2 \theta_W)] F_{q\bar{q}}(s, Y) \quad (6)$$

$$\sin^2 \theta_W = 0.2287 \pm 0.0020(stat.) \pm 0.0025(syst.) \quad (7)$$

Systematic uncertainties in the measurement of $\sin^2 \theta_W$:

| Source | Correction | Uncertainty |
|---------------------------|------------|--------------|
| PDF | - | ± 0.0013 |
| FSR | - | ± 0.0011 |
| LO model (EWK) | - | ± 0.0002 |
| LO model (QCD) | +0.0012 | ± 0.0012 |
| Resolution/alignment | +0.0007 | ± 0.0013 |
| Acceptance and Efficiency | - | ± 0.0003 |
| Background | - | ± 0.0001 |
| Total | +0.0019 | ± 0.0025 |

Phys. Rev. D 84, 112002 (2011)

- Drell-Yan process is a good probe for new physics
- $\sin^2 \theta_W$, A_{FB} and DY cross section measurements are consistent with the Standard Model predictions within uncertainties