

Extended scalar sector

- 2HDM Type II Yukawa
- IDM2

QFTHEP
June 2013

Per Osland

Work with:

L Basso, A Lipniacka, F Mahmoudi, S Moretti, G M Pruna, M Purmohammadi

G M Pruna, A Pukhov, M Purmohammadi

arXiv:1205.6569, arXiv:1305.3219

Preamble

- Higgs particle found! SM?
- 2HDM excluded?
- not quite
- but parameter space severely constrained
- Look for charged Higgs!

2HDM notation 1

$$\begin{aligned} V = & \frac{\lambda_1}{2} (\Phi_1^\dagger \Phi_1)^2 + \frac{\lambda_2}{2} (\Phi_2^\dagger \Phi_2)^2 + \lambda_3 (\Phi_1^\dagger \Phi_1) (\Phi_2^\dagger \Phi_2) \\ & + \lambda_4 (\Phi_1^\dagger \Phi_2) (\Phi_2^\dagger \Phi_1) + \frac{1}{2} \left[\lambda_5 (\Phi_1^\dagger \Phi_2)^2 + \text{h.c.} \right] \\ & - \frac{1}{2} \left\{ m_{11}^2 (\Phi_1^\dagger \Phi_1) + \left[m_{12}^2 (\Phi_1^\dagger \Phi_2) + \text{h.c.} \right] + m_{22}^2 (\Phi_2^\dagger \Phi_2) \right\} \end{aligned}$$

No FCNC:

$$\lambda_6 = 0; \quad \lambda_7 = 0$$

Allow CPV: $\lambda_5, \quad m_{12}^2 \quad \text{complex}$

2HDM notation 2

$$\Phi_i = \begin{pmatrix} \varphi_i^+ \\ \frac{1}{\sqrt{2}}(v_i + \eta_i + i\chi_i) \end{pmatrix}$$

$$\begin{pmatrix} H_1 \\ H_2 \\ H_3 \end{pmatrix} = R \begin{pmatrix} \eta_1 \\ \eta_2 \\ \eta_3 \end{pmatrix}$$

$$\eta_3 = -\sin \beta \chi_1 + \cos \beta \chi_2$$

$$R\mathcal{M}^2 R^T = \mathcal{M}_{\text{diag}}^2 = \text{diag}(M_1^2, M_2^2, M_3^2)$$

2HDM notation 3

$$\begin{aligned}
 & \qquad \qquad \qquad 2 \text{ vs } 3 \qquad \qquad \qquad 1 \text{ vs } 3 \qquad \qquad \qquad 1 \text{ vs } 2 \\
 R = R_3 R_2 R_1 &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \alpha_3 & \sin \alpha_3 \\ 0 & -\sin \alpha_3 & \cos \alpha_3 \end{pmatrix} \begin{pmatrix} \cos \alpha_2 & 0 & \sin \alpha_2 \\ 0 & 1 & 0 \\ -\sin \alpha_2 & 0 & \cos \alpha_2 \end{pmatrix} \begin{pmatrix} \cos \alpha_1 & \sin \alpha_1 & 0 \\ -\sin \alpha_1 & \cos \alpha_1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\
 &= \begin{pmatrix} c_1 c_2 & s_1 c_2 & s_2 \\ -(c_1 s_2 s_3 + s_1 c_3) & c_1 c_3 - s_1 s_2 s_3 & c_2 s_3 \\ -c_1 s_2 c_3 + s_1 s_3 & -(c_1 s_3 + s_1 s_2 c_3) & c_2 c_3 \end{pmatrix} \quad \text{PDG convention}
 \end{aligned}$$

$$c_i = \cos \alpha_i, \quad s_i = \sin \alpha_i$$

CP-conserving limits:

H_1 odd: $\alpha_2 \simeq \pm\pi/2$, α_1, α_3 arbitrary,

H_2 odd: $\alpha_2 = 0$, $\alpha_3 = \pi/2$, α_1 arbitrary,

H_3 odd: $\alpha_2 = \alpha_3 = 0$, α_1 arbitrary.

Yukawa couplings

$$H_j b \bar{b} : \quad \frac{-ig m_b}{2 m_W} \frac{1}{\cos \beta} [R_{j1} - i\gamma_5 \sin \beta R_{j3}],$$

$$H_j t \bar{t} : \quad \frac{-ig m_t}{2 m_W} \frac{1}{\sin \beta} [R_{j2} - i\gamma_5 \cos \beta R_{j3}].$$

$$H^+ b \bar{t} : \quad \frac{ig}{2\sqrt{2} m_W} V_{tb} [m_b(1 + \gamma_5) \tan \beta + m_t(1 - \gamma_5) \cot \beta],$$

$$H^- t \bar{b} : \quad \frac{ig}{2\sqrt{2} m_W} V_{tb}^* [m_b(1 - \gamma_5) \tan \beta + m_t(1 + \gamma_5) \cot \beta].$$

Gauge couplings

$$H_j Z Z : \quad [\cos \beta R_{j1} + \sin \beta R_{j2}], \quad \text{for } j = 1,$$

$$H_1 \rightarrow Z Z, W W \quad \text{observed (off-shell)}$$

$$H_{2,3} \rightarrow Z Z, W W \quad \text{not observed (on-shell)}$$

$$H_j H^\pm W^\mp : \quad \frac{g}{2} [\mp i (\sin \beta R_{j1} - \cos \beta R_{j2}) + R_{j3}] (p_\mu^j - p_\mu^\pm).$$

Enter in total widths: $H_{2,3} \rightarrow H_1 Z$

Parameters

Input: $\tan \beta, (M_1, M_2), (M_{H^\pm}, \mu^2), (\alpha_1, \alpha_2, \alpha_3)$

Reconstruct:


$$M_3^2 = \frac{M_1^2 R_{13}(R_{12} \tan \beta - R_{11}) + M_2^2 R_{23}(R_{22} \tan \beta - R_{21})}{R_{33}(R_{31} - R_{32} \tan \beta)}$$

Explicit expressions for

$$\lambda_1, \lambda_2, \lambda_3, \lambda_4, \text{Re } \lambda_5, \text{Im } \lambda_5$$

in terms of input


Branching ratios



Branching ratios

P4

2HDM: H_1 branching ratios




Decay rates

random

2HDM: Decay rate

$\tan\beta = 1$




Decay rates

P4

2HDM: Decay rate

$\tan\beta = 1$




Decay rates

random

2HDM: Decay rate

$\tan\beta = 2$




Decay rates

P8


2HDM: Decay rate

$\tan\beta = 2$



SM decay rate

has stronger coupling to WW



H_2 (if, for example at 400 GeV) and H_3
must decay more slowly than SM Higgs (at same mass),
in order for model not to be excluded by LHC data

Constraints-theory

- Positivity
 - Explicit conditions
- Unitarity
 - Explicit conditions
- Perturbativity
- Global minimum
 - Three coupled cubic equations

Constraints-experiment

- $b \rightarrow s\gamma$
- $\Gamma(Z \rightarrow b\bar{b})$
- $B \rightarrow \tau\nu(X), B \rightarrow D\tau\nu, D \rightarrow \tau\nu$
- $B_0 \leftrightarrow \bar{B}_0$
- $B_{d,s} \rightarrow \mu^+\mu^-$
- EW constraints: S, T
- Electron EDM
- LHC: $H_1 \rightarrow \gamma\gamma$
- LHC: $H_{2,3} \rightarrow W^+W^-$

Parameters

Input:

$\tan \beta, (M_1, M_2), (M_{H^\pm}, \mu^2), (\alpha_1, \alpha_2, \alpha_3)$



Typically: step



fix

step

scan

Allowed regions (red)

Ignore LHC (apologies)



LHC constraints

1 $gg \rightarrow H_1 \rightarrow \gamma\gamma$

$$R_{\gamma\gamma} = \frac{\Gamma(H_1 \rightarrow gg)\text{BR}(H_1 \rightarrow \gamma\gamma)}{\Gamma(H_{\text{SM}} \rightarrow gg)\text{BR}(H_{\text{SM}} \rightarrow \gamma\gamma)}$$

Triangle diagrams modified by couplings, also axial term

$$0.5 \leq R_{\gamma\gamma} \leq 2.0$$


2 $gg \rightarrow H_{2,3} \rightarrow W^+W^-$

$$R_{ZZ} = \frac{\Gamma(H_j \rightarrow gg)\text{BR}(H_j \rightarrow ZZ)}{\Gamma(H_{\text{SM}} \rightarrow gg)\text{BR}(H_{\text{SM}} \rightarrow ZZ)}$$


bounded

Adopt LHC (ATLAS & CMS) 95% CL

ATLAS CMS





ATLAS, CMS, CMS 2013



Allowed regions





LHC constraints 2012



Next:

- **Combine all constraints:**





Allowed regions (green) 2012







2013 vs 2012

- 2013: More exclusion in heavy-Higgs region
- 2012 $0.5 \leq R_{\gamma\gamma} \leq 2.0$
- 2013 ATLAS $1.03 \leq R_{\gamma\gamma} \leq 2.33 \quad (2\sigma)$
- 2013 CMS $0.26 \leq R_{\gamma\gamma} \leq 1.34 \quad (2\sigma)$





Allowed regions 2013



Allowed regions 2013



Allowed regions 2013



$$\tan \beta = 1$$

Range of μ (all masses in GeV)

$M_{H^\pm} \backslash M_2$	300	350	400	450	500
500	none	none	(300,400)	(300,450)	(350,500]
450	none	[250,350]	(250,400]	(250,450]	(300,450]
400	(300)	(200,350]	(200,400]	(250,400]	(300,400]

$$\mu \lesssim \text{avg}(M_2, M_{H^\pm})$$





$$\tan \beta = 2$$

Range of μ (all masses in GeV)

$M_{H^\pm} \backslash M_2$	300	350	400	450	500
500	none	none	(400,450)	(400,450)	(400,500]
450	none	(300,350)	(300,450)	(350,450]	(400,500]
400	(200,300]	(200,350]	(250,450]	(300,450]	(350,400]



$$\mu \lesssim \text{avg}(M_2, M_{H^\pm})$$

Allowed regions high mass



Only CP-conserving limits survive
at high masses

Allowed regions high tanbeta 2012



Decoupling



$A=H_2, A=H_3$

Decoupling 1: $(\alpha_1, \alpha_2) \sim (\pm\pi/2, 0)$


Decoupling 2: $(\alpha_1, \alpha_2) \sim (0, \pm\pi/2)$

$A=H_1$ Excluded by LHC



Allowed regions high tanbeta 2013





Allowed regions high tanbeta 2013



CP conserving limits





CP conserving limits





$$H_1 \rightarrow \gamma\gamma$$

- $R_{\gamma\gamma} > 1$?
- In SM W and t loop interfere destructively
- $H_j t\bar{t}$: $\frac{-ig m_t}{2 m_W} \frac{1}{\sin \beta} [R_{j2} - i\gamma_5 \cos \beta R_{j3}]$.
- Flip sign of t -loop?
- $R_{12} = s_1 c_2$, $s_1 < 0?$ $c_2 < 0?$
- Also γ_5 term (additive)

Overview ATLAS constraints




Hermann et al
2012
(lower bound)




Overview

CMS constraints



H_3 mass, M_3 CMS constraints



Charged Higgs

- Only way to exclude SM?
- Identify Benchmarks!

Requirements:

- Not excluded by theoretical arguments
- Not excluded by experimental data
- Good production cross section
- Good BR for decay to $W + H_1$
- Moderate background

Charged Higgs Benchmarks

	α_1/π	α_2/π	α_3/π	$\tan \beta$	M_2	$M_{H^\pm}^{\min}, M_{H^\pm}^{\max}$
P_1	0.23	0.06	0.005	1	300	300,325
P_2	0.35	-0.014	0.48	1	300	300,415
P_3	0.35	-0.015	0.496	1	350	300,450
P_4	0.35	-0.056	0.43	1	400	300,455
P_5	0.33	-0.21	0.23	1	450	300,470
P_6	0.27	-0.26	0.25	1	500	300,340
P_7	0.39	-0.07	0.33	2	300	300,405
P_8	0.34	-0.03	0.11	2	400	300,315
P_9	0.47	-0.006	0.05	10	400	400,440
P_{10}	0.49	-0.002	0.06	10	600	600,700

Proposed channel:

$$pp \rightarrow W^\pm H^\mp (+X)$$

$$\rightarrow W^+ W^- H_1$$

$$\rightarrow \underbrace{jj}_W \underbrace{\ell^\pm \nu}_W \underbrace{b\bar{b}}_{H_1}$$

Proposed channel:

$$pp \rightarrow W^\pm H^\mp (+X)$$

$$\rightarrow W^+ W^- H_1$$

$$\rightarrow \underbrace{jj}_W \underbrace{\ell^\pm \nu}_W \underbrace{b\bar{b}}_{H_1}$$

$H_j H^\pm W^\mp$ coupling squared:

$$\sim (\sin \beta R_{j1} - \cos \beta R_{j2})^2 + R_{j3}^2$$

$$H_1 H^\pm W^\mp : = \sin^2(\beta - \alpha_1) \cos^2 \alpha_2 + \sin^2 \alpha_2$$

Proposed channel:

$$pp \rightarrow W^\pm H^\mp (+X)$$

$$\rightarrow W^+ W^- H_1$$



$$\rightarrow \underbrace{jj}_W \underbrace{\ell^\pm \nu}_W \underbrace{b\bar{b}}_{H_1}$$

$H_j H^\pm W^\mp$ coupling squared:



$$\sim (\sin \beta R_{j1} - \cos \beta R_{j2})^2 + R_{j3}^2$$

$$H_1 H^\pm W^\mp : = \sin^2(\beta - \alpha_1) + \sin^2 \alpha_2 \cos^2(\beta - \alpha_1)$$

Branching ratios:




Branching ratios:





Dominant production mechanisms



Coupling may depend on details





(a) irreducible background




Cross sections: legend next page



Cross sections:



Background

- $t\bar{t} \rightarrow b\bar{b}W^+W^-$ 
- cross section larger by factor 10^3
- impose generic cuts, BG reduction by factor 40, signal reduction by 2-3

Generic cuts

1) **Kinematics:** standard detector cuts

$$\begin{aligned} p_\ell^T &> 15 \text{ GeV}, & |\eta_\ell| &< 2.5, \\ p_j^T &> 20 \text{ GeV}, & |\eta_j| &< 3, \\ |\Delta R_{jj}| &> 0.5, & |\Delta R_{\ell j}| &> 0.5; \end{aligned}$$

2) **light Higgs reconstruction:**

$$|M(b\bar{b}) - 125 \text{ GeV}| < 20 \text{ GeV};$$

3) **hadronic W reconstruction ($W_h \rightarrow jj$):**

$$|M(jj) - 80 \text{ GeV}| < 20 \text{ GeV};$$

Generic cuts

4) **top veto:** if $\Delta R(b_1, W_h) < \Delta R(b_2, W_h)$, then

$$M(b_1 jj) > 200 \text{ GeV}, \quad M_T(b_2 \ell \nu) > 200 \text{ GeV},$$

otherwise $1 \leftrightarrow 2$; **disfavor top, for each b-quark separately**


5) **same-hemisphere b quarks:**

$$\frac{\mathbf{p}_{b_1}}{|\mathbf{p}_{b_1}|} \cdot \frac{\mathbf{p}_{b_2}}{|\mathbf{p}_{b_2}|} > 0.$$

Additional anti-top cut

Idea: Since $M_{H^\pm} > m_t$

One of the W's should form
high invariant mass with $b\bar{b}$ pair



Possible cuts

“squared cut”: $C_{\text{squ}} = \max (M(b\bar{b}jj), M_T(b\bar{b}l\nu)) > M_{\text{lim}}$



“single cut”: $C_{\text{sng}} = M_T(b\bar{b}l\nu) > M_{\text{lim}} .$

Choose:



C_{sng}

$M_{\text{lim}} = 600 \text{ GeV}$



$$P_2 : \tan \beta = 1, \quad M_2 = 300 \text{ GeV}, \quad \alpha_i = \{0.35, -0.014, 0.48\}$$





$$P_3 : \tan \beta = 1, \quad M_2 = 350 \text{ GeV}, \quad \alpha_i = \{0.35, -0.015, 0.496\}$$





$$P_4 : \tan \beta = 1, \quad M_2 = 400 \text{ GeV}, \quad \alpha_i = \{0.35, -0.056, 0.43\}$$



$$P_5 : \tan \beta = 1, \quad M_2 = 450 \text{ GeV}, \quad \alpha_i = \{0.33, -0.21, 0.23\}$$



$$P_7 : \tan \beta = 2, \quad M_2 = 300 \text{ GeV}, \quad \alpha_i = \{0.39, -0.07, 0.33\}$$



	$M_{H^\pm} = 310 \text{ GeV}$		$M_{H^\pm} = 390 \text{ GeV}$	
	Events	S/\sqrt{B}	Events	S/\sqrt{B}
$t\bar{t}$	24.9			
peak	11.9	—	9.9	—
P_1	3.8	0.8	—	—
peak	2.6	0.8	—	—
P_2	4.7	1.0	8.8	1.8
peak	3.3	1.0	7.3	2.3
P_3	11.3	2.3	22.0	4.4
peak	7.7	2.3	17.2	5.4
P_4	10.0	2.0	20.3	4.1
peak	7.8	2.3	16.0	5.1
P_5	21.1	4.2	30.2	6.1
peak	13.9	4.1	25.0	7.9
P_6	14.0	2.8	—	—
peak	9.4	2.8	—	—
P_7	3.1	0.6	7.4	1.5
peak	2.8	0.8	7.3	2.3
P_8	1.2	0.2	—	—
peak	1.2	0.4	—	—

2HDM Conclusions

- 2HDM II parameter space is severely constrained by LHC data
- Parts of 2HDM II parameter space are still open
- SM would be excluded by charged Higgs discovery
- $pp \rightarrow \underbrace{jj}_W \underbrace{\ell^\pm \nu}_W \underbrace{b\bar{b}}_{H_1}$ channel allows detection in part of parameter space

The Extension

Scalar DM

- “Inert (Scalar) Doublet Model (IDM)”, Barbieri et al, 2006
Extend SM with additional scalar doublet, unbroken Z_2 symmetry makes lightest “odd” particle stable. No vev, no direct coupling to SM matter.
- “CP-violating Inert Doublet Model”, Grzadkowski et al, 2009
Extend 2HDM with additional scalar doublet, unbroken Z_2 symmetry makes lightest “odd” particle stable. No vev, no direct coupling to SM matter.

IDM2: 2HDM + inert doublet

Grzadkowski et al, 2009

Motivation: IDM + CP violation

Fields:

$$\Phi_1 = \begin{pmatrix} \varphi_1^+ \\ (v_1 + \eta_1 + i\chi_1)/\sqrt{2} \end{pmatrix}, \quad \Phi_2 = \begin{pmatrix} \varphi_2^+ \\ (v_2 + \eta_2 + i\chi_2)/\sqrt{2} \end{pmatrix}$$

$$\eta = \begin{pmatrix} \eta^+ \\ (S + iA)/\sqrt{2} \end{pmatrix}$$

Potential:

$$V(\Phi_1, \Phi_2, \eta) = V_{12}(\Phi_1, \Phi_2) + V_3(\eta) + V_{123}(\Phi_1, \Phi_2, \eta)$$

Coupling:



$$V_{12}(\Phi_1, \Phi_2) = -\frac{1}{2} \left\{ m_{11}^2 \Phi_1^\dagger \Phi_1 + m_{22}^2 \Phi_2^\dagger \Phi_2 + \left[m_{12}^2 \Phi_1^\dagger \Phi_2 + \text{h.c.} \right] \right\}$$

(standard)

$$+ \frac{\lambda_1}{2} (\Phi_1^\dagger \Phi_1)^2 + \frac{\lambda_2}{2} (\Phi_2^\dagger \Phi_2)^2 + \lambda_3 (\Phi_1^\dagger \Phi_1) (\Phi_2^\dagger \Phi_2)$$

$$+ \lambda_4 (\Phi_1^\dagger \Phi_2) (\Phi_2^\dagger \Phi_1) + \frac{1}{2} \left[\lambda_5 (\Phi_1^\dagger \Phi_2)^2 + \text{h.c.} \right]$$

$$V_3(\eta) = m_\eta^2 \eta^\dagger \eta + \frac{\lambda_\eta}{2} (\eta^\dagger \eta)^2$$

Coupling:

(most general)

$$V_{123}(\Phi_1, \Phi_2, \eta) = \lambda_{1133} (\Phi_1^\dagger \Phi_1) (\eta^\dagger \eta) + \lambda_{2233} (\Phi_2^\dagger \Phi_2) (\eta^\dagger \eta)$$

$$+ \lambda_{1331} (\Phi_1^\dagger \eta) (\eta^\dagger \Phi_1) + \lambda_{2332} (\Phi_2^\dagger \eta) (\eta^\dagger \Phi_2)$$

$$+ \frac{1}{2} \left[\lambda_{1313} (\Phi_1^\dagger \eta)^2 + \text{h.c.} \right] + \frac{1}{2} \left[\lambda_{2323} (\Phi_2^\dagger \eta)^2 + \text{h.c.} \right]$$

Many parameters...

Many parameters! Simplify!

“Dark democracy”:

$$\lambda_a \equiv \lambda_{1133} = \lambda_{2233},$$
$$\lambda_b \equiv \lambda_{1331} = \lambda_{2332},$$
$$\lambda_c \equiv \lambda_{1313} = \lambda_{2323} \text{ (real).}$$

Masses of inert sector:

$$M_{\eta^\pm}^2 = m_\eta^2 + \frac{1}{2}\lambda_a v^2,$$
$$M_S^2 = m_\eta^2 + \frac{1}{2}(\lambda_a + \lambda_b + \lambda_c)v^2 = M_{\eta^\pm}^2 + \frac{1}{2}(\lambda_b + \lambda_c)v^2,$$
$$M_A^2 = m_\eta^2 + \frac{1}{2}(\lambda_a + \lambda_b - \lambda_c)v^2 = M_{\eta^\pm}^2 + \frac{1}{2}(\lambda_b - \lambda_c)v^2$$

Important:

These $\lambda_{a,b,c}$ characterize **coupling** of inert sector to non-inert sector, and also **mass splitting** in inert sector

Higgs portal

- Coupling of scalars: Higgs \longleftrightarrow DM

$$\lambda_L \equiv \frac{1}{2}(\lambda_a + \lambda_b + \lambda_c) = \frac{M_S^2 - m_\eta^2}{v^2}$$

Constraints

- positivity (rather complicated), 20% excluded
- unitarity, 60% excluded
- global minimum, 10% excluded
- additional 2HDM constraints: T , $b \rightarrow s\gamma$ etc
- DM

determined by MicrOMEGAs



EW “precision data”



Positivity

Define:

$$\lambda_x = \lambda_3 + \min(0, \lambda_4 - |\lambda_5|)$$

$$\lambda_y = \lambda_{1133} + \min(0, \lambda_{1331} - |\lambda_{1313}|)$$

$$\lambda_z = \lambda_{2233} + \min(0, \lambda_{2332} - |\lambda_{2323}|)$$

$$\lambda_1 > 0, \quad \lambda_2 > 0, \quad \lambda_\eta > 0, \quad \lambda_x > -\sqrt{\lambda_1 \lambda_2}$$

$$\lambda_y > -\sqrt{\lambda_1 \lambda_\eta}, \quad \lambda_z > -\sqrt{\lambda_2 \lambda_\eta}$$

Plus additional constraint, which in the case of
Dark democracy $\lambda_y = \lambda_z$ takes the form:

$$\lambda_y \geq 0 \vee \left(\lambda_\eta \lambda_x - \lambda_y^2 > -\sqrt{(\lambda_\eta \lambda_1 - \lambda_y^2)(\lambda_\eta \lambda_2 - \lambda_y^2)} \right)$$

Getting correct DM density

Main Early Universe annihilation mechanisms:

- Annihilation to $W^+ W^-$, effective above 75 GeV
- Annihilation via real or virtual neutral Higgs

like IDM...

Annihilation in the Early Universe

The DM particles can annihilate via the gauge coupling:

$$SSW^+W^- : \frac{ig^2}{2}$$
$$SSZZ : \frac{ig^2}{2\cos^2\theta_W}$$

Annihilation in the Early Universe

The DM particles can annihilate via the gauge coupling:

$$SSW^+W^- : \frac{ig^2}{2}$$

$$SSZZ : \frac{ig^2}{2\cos^2\theta_W}$$

or to non-inert scalars via trilinear

$$SSH_j : -2iF_{SSj}\lambda_L v$$

$$F_{SSj} = \cos\beta R_{j1} + \sin\beta R_{j2}$$

or quartic couplings:

$$SSH_j H_j : -2i(\lambda_L - \lambda_c R_{j3}^2)$$



$$SSH_j H_k : 2i\lambda_c R_{j3} R_{k3}$$

$$SSH^+ H^- : -i\lambda_a$$

Note



Allowed regions in M_s




Scan over parameters

1. M_S, M_1 (lowest masses of inert and 2HDM sectors, fixed)
2. M_A, M_{η^\pm} (inert sector, physical masses, fixed).
3. M_2, μ (2HDM sector parameters)
4. m_η (inert sector, soft mass parameter, fixed).
5. $\tan \beta, M_{H^\pm}$ (2HDM sector),
 $0.5 \leq \tan \beta \leq 50, 300 \text{ GeV} \leq M_{H^\pm} \leq 700 \text{ GeV}.$
6. $\alpha_1, \alpha_2, \alpha_3$ (2HDM sector),
 $-\pi/2 \leq \alpha_{1,2} \leq \pi/2, \text{ and } 0 \leq \alpha_3 \leq \pi/2.$

Collect results in M_{η^\pm}, m_η plane

Need some coupling to Higgs, $|\lambda_L|$ can not be too small




Mass of charged state

Compare CDMS-II and XENON100

2011


Cross sections



Compare CDMS-II and XENON100


2011

2013




Can the model be experimentally tested?

Production and discovery at the LHC?




Pair production and single production



Decay:





Branching ratios of η^\pm to A and S.



Single production




Pair production




Bumps are due to resonant production via H_2, H_3

For small mass splitting, get displaced vertex



Decay:







For hadronic W decay, get two jets (may merge to one)

Ideas for a search

$$pp \rightarrow \eta^\pm S \rightarrow W^\pm SS.$$

$$pp \rightarrow j + \text{MET}$$

$$\text{MET} > 120 \text{ GeV}, \quad p_j^T > 120 \text{ GeV}$$



Conclusions

...if scalars are dark matter...

- Scalar sector could be much more exciting than in the SM
- Possibly signals in Direct or Indirect detection experiments
- Possibly interesting signals at the LHC
- In the meantime, parts of parameter space will be excluded