

The Feynman-Vernon Influence Functional Approach in QED


Mark Shleenkov, Alexander Biryukov

Samara State University
General and Theoretical Physics Department

The XXII International Workshop
High Energy Physics and Quantum Field Theory
June 24 — July 1, 2015
Samara, Russia

The Feynman-Vernon Influence Functional approach

R.P. Feynman, F.L. Vernon, Jr. **The Theory of a General Quantum System Interacting with a Linear Dissipative System**,
Annals of Physics **24**, 118–173.



General quantum systems Q and X coupled by a potential $V(Q, X, t)$.

«... It is shown that the effect of the external systems in such a formalism [paths integral formalism] can always be included in a general class of functionals (influence functionals) of the coordinates of the system only ... »

QED Lagrangian

$$\mathcal{L}(x) = \bar{\psi}(x)(i\gamma^\mu \partial_\mu - m)\psi(x) - \frac{1}{4}F_{\mu\nu}(x)F^{\mu\nu}(x) - ej^\mu(x)A_\mu(x) \quad (1)$$

where

$$F_{\mu\nu}(x) = \partial_\mu A_\nu(x) - \partial_\nu A_\mu(x) \quad (2)$$

$$j^\mu(x) = \bar{\psi}(x)\gamma^\mu\psi(x) \quad (3)$$

$$\hat{\psi}(\mathbf{x}, t) = \sum_{\mathbf{p}, \sigma=1,2} \frac{1}{\sqrt{2V\omega_{\mathbf{p}}^{(f)}}} \left(\hat{b}_{\mathbf{p}\sigma} u_{\sigma}(p) e^{ipx} + \hat{c}_{\mathbf{p}\sigma}^{\dagger} u_{\sigma}(-p) e^{-ipx} \right), \quad (4)$$

$$\hat{\bar{\psi}}(\mathbf{x}, t) = \sum_{\mathbf{p}, \sigma=1,2} \frac{1}{\sqrt{2V\omega_{\mathbf{p}}^{(f)}}} \left(\hat{b}_{\mathbf{p}\sigma}^{\dagger} \bar{u}_{\sigma}(p) e^{-ipx} + \hat{c}_{\mathbf{p}\sigma} \bar{u}_{\sigma}(-p) e^{ipx} \right), \quad (5)$$

$$\hat{j}_{\mu}(\mathbf{x}, t) = \hat{\bar{\psi}}(\mathbf{x}, t) \gamma_{\mu} \hat{\psi}(\mathbf{x}, t) \quad (6)$$

$$\hat{A}^{\mu}(\mathbf{x}, t) = \sum_{\mathbf{k}, \lambda=1,2} \frac{1}{\sqrt{2V\omega_{\mathbf{k}}^{(b)}}} \varepsilon_{\lambda}^{\mu} \left(\hat{a}_{\mathbf{k}\lambda} e^{ikx} + \hat{a}_{\mathbf{k}\lambda}^{\dagger} e^{-ikx} \right) \quad (7)$$

Second quantization

$$\hat{H}_{full} = \sum_{\mathbf{p}, \sigma=1,2} \omega_{\mathbf{p}}^{(f)} \left(\hat{b}_{\mathbf{p}\sigma}^\dagger \hat{b}_{\mathbf{p}\sigma} + \hat{c}_{\mathbf{p}\sigma}^\dagger \hat{c}_{\mathbf{p}\sigma} \right) + \sum_{\mathbf{k}, \lambda=1,2} \omega_{\mathbf{k}}^{(b)} \hat{a}_{\mathbf{k}\lambda}^\dagger \hat{a}_{\mathbf{k}\lambda} + e \sum_{\mathbf{k}, \lambda=1,2} \frac{\imath}{\sqrt{2\omega_{\mathbf{k}}^{(b)} V}} \left(\varepsilon_\lambda^\mu \hat{j}_\mu^+(\mathbf{k}, t) \hat{a}_{\mathbf{k}\lambda} + \varepsilon_\lambda^{*\mu} \hat{j}_\mu^-(\mathbf{k}, t) \hat{a}_{\mathbf{k}\lambda}^\dagger \right) \quad (8)$$

where

$$\hat{j}_\mu^+(\mathbf{k}, t) = \int \hat{j}_\mu(\mathbf{x}, t) e^{\imath \mathbf{kx}} d\mathbf{x}, \quad \hat{j}_\mu^-(\mathbf{k}, t) = \int \hat{j}_\mu(\mathbf{x}, t) e^{-\imath \mathbf{kx}} d\mathbf{x} \quad (9)$$

Evolution equation for statistical operator $\hat{\rho}(t_f)$

$$\hat{\rho}(t_f) = \hat{U}(t_f, t_{in})\hat{\rho}(t_{in})\hat{U}^\dagger(t_f, t_{in}) \quad (10)$$

where $\hat{\rho}(t_{in})$ is statistical operator, describing initial state at moment t_{in} ,
 $\hat{U}(t_f, t_{in})$ — evolution operator.

$$\hat{U}(t_f, t_{in}) = \hat{T} \exp\left[-\frac{i}{\hbar} \int_{t_{in}}^{t_f} \hat{H}_{full}(\tau) d\tau\right]. \quad (11)$$

where \hat{H}_{full} :

$$\hat{H}_{full} = \hat{H}_{sys} + \hat{H}_{field} + \hat{H}_{int} \quad (12)$$

Evolution equation for density matrix in holomorphic representation

$$|\theta_{p\lambda}, \alpha_{k\lambda}\rangle = |\theta_{p\lambda}\rangle \otimes |\alpha_{k\lambda}\rangle$$

The density matrix:

$$\rho(\alpha_f^*, \bar{\theta}_f, \alpha'_f, \theta'_f; t_f) = \langle \bar{\theta}_f, \alpha_f | \hat{\rho}(t_f) | \theta'_f, \alpha'_f \rangle \quad (13)$$

The kernel of evolution operator:

$$U(\alpha_f^*, \bar{\theta}_f, t_f | \alpha_{in}, \theta_{in}, t_{in}) = \langle \bar{\theta}_f, \alpha_f | \hat{U}(t_f, t_{in}) | \theta_{in}, \alpha_{in} \rangle \quad (14)$$

The evolution equation:

$$\begin{aligned} \rho(\alpha_f^*, \bar{\theta}_f, \alpha'_f, \theta'_f; t_f) &= \int \frac{d^2 \alpha'_{in}}{\pi} \frac{d^2 \theta'_{in}}{\pi} \frac{d^2 \alpha_{in}}{\pi} \frac{d^2 \theta_{in}}{\pi} \times \\ &\times U(\alpha_f^*, \bar{\theta}_f, t_f | \alpha_{in}, \theta_{in}, t_{in}) \rho(\alpha_{in}^*, \bar{\theta}_{in}, \alpha'_{in}, \theta'_{in}; t_{in}) U^*(\alpha'_f, \theta'_f, t_f | \alpha'^*_{in}, \bar{\theta}'_{in}; t_{in}) \end{aligned} \quad (15)$$

Coherent states for electromagnetic field

$$\hat{a}_{\mathbf{k}\lambda}|\alpha_{\mathbf{k}\lambda}\rangle = \alpha_{\mathbf{k}\lambda}|\alpha_{\mathbf{k}\lambda}\rangle, \quad \langle\alpha_{\mathbf{k}\lambda}|\hat{a}_{\mathbf{k}\lambda}^\dagger = \langle\alpha_{\mathbf{k}\lambda}|\alpha_{\mathbf{k}\lambda}^*, \quad (16)$$

where $\alpha_{\mathbf{k}\lambda}$ — complex value, which describe states \mathbf{k} mode of quantum electromagnetic field. These states ($|\alpha\rangle$) are non-orthogonal:

$$\langle\alpha'_{\mathbf{k}'\lambda'}|\alpha_{\mathbf{k}\lambda}\rangle = \delta_{\mathbf{k}'\mathbf{k}}\delta_{\lambda'\lambda} \exp\left\{-\frac{1}{2}\left(|\alpha'_{\mathbf{k}'\lambda'}|^2 + |\alpha_{\mathbf{k}\lambda}|^2 - 2\alpha'^*_{\mathbf{k}'\lambda'}\alpha_{\mathbf{k}\lambda}\right)\right\}. \quad (17)$$

There is resolution of the identity operator:

$$\int |\alpha_{\mathbf{k}\lambda}\rangle\langle\alpha_{\mathbf{k}\lambda}| \frac{d^2\alpha_{\mathbf{k}\lambda}}{\pi} = \hat{1}. \quad (18)$$

Grassmann states for Dirac field

$$\hat{b}_{\mathbf{p},\sigma} |\theta_{\mathbf{p},\sigma}\rangle = \theta_{\mathbf{p},\sigma} |\theta_{\mathbf{p},\sigma}\rangle, \quad \langle \bar{\theta}_{\mathbf{p},\sigma} | \hat{b}_{\mathbf{p},\sigma}^\dagger = \langle \bar{\theta}_{\mathbf{p},\sigma} | \bar{\theta}_{\mathbf{p},\sigma}, \quad (19)$$

where $\theta_{\mathbf{p},\sigma}$ — grassman variable. These states ($|\theta\rangle$) are non-orthogonal:

$$\langle \bar{\theta}'_{\mathbf{p}'\sigma'} | \theta_{\mathbf{p}\sigma} \rangle = \delta_{\mathbf{p}'\mathbf{p}} \delta_{\sigma'\sigma} \exp \left\{ -\frac{1}{2} \left(\bar{\theta}'_{\mathbf{p}'\sigma'} \theta'_{\mathbf{p}'\sigma'} + \bar{\theta}_{\mathbf{p}\sigma} \theta_{\mathbf{p}\sigma} - 2 \bar{\theta}'_{\mathbf{p}'\sigma'} \theta_{\mathbf{p}\sigma} \right) \right\}. \quad (20)$$

There is resolution of the identity operator:

$$\int |\theta_{\mathbf{p}\sigma}\rangle \langle \bar{\theta}_{\mathbf{p}\sigma}| \frac{d^2\theta_{\mathbf{p}\sigma}}{\pi} = \hat{1}. \quad (21)$$

The kernel of evolution operator

$$U(\alpha_f^*, \bar{\theta}_f, t_f | \alpha_f, \theta_f, t_{in}) = \int \mathfrak{D}\alpha^*(\tau) \mathfrak{D}\alpha(\tau) \mathfrak{D}\bar{\theta}(\tau) \mathfrak{D}\theta(\tau) \times \\ \times \exp \left\{ iS_{full} [\alpha^*(\tau), \alpha(\tau), \bar{\theta}(\tau), \theta(\tau)] \right\}, \quad (22)$$

where action

$$S_{full} [\alpha^*(\tau), \alpha(\tau), \bar{\theta}(\tau), \theta(\tau)] = \\ = S_f [\bar{\theta}(\tau), \theta(\tau)] + S_b [\alpha^*(\tau), \alpha(\tau)] + S_{int} [\alpha^*(\tau), \alpha(\tau), \bar{\theta}(\tau), \theta(\tau)]. \quad (23)$$

Action of fermionic field:

$$S_f [\bar{\theta}(\tau), \theta(\tau)] = \int_{t_{in}}^{t_f} \left(\frac{\dot{\bar{\theta}}(\tau)\theta(\tau) - \bar{\theta}(\tau)\dot{\theta}(\tau)}{2i} - \omega^{(f)}\bar{\theta}(\tau)\theta(\tau) \right) d\tau \quad (24)$$

Action of bosonic field:

$$S_b [\alpha^*(\tau), \alpha(\tau)] = \int_{t_{in}}^{t_f} \left(\frac{\dot{\alpha}^*(\tau)\alpha(\tau) - \alpha^*(\tau)\dot{\alpha}(\tau)}{2i} - \omega^{(b)}\alpha^*(\tau)\alpha(\tau) \right) d\tau \quad (25)$$

Action of interaction part:

$$S_{int} [\alpha^*(\tau), \alpha(\tau), \bar{\theta}(\tau), \theta(\tau)] = \int_{t_{in}}^{t_f} ej^\mu(\bar{\theta}(\tau), \theta(\tau))(\varepsilon_\mu^*\alpha^*(\tau) + \varepsilon_\mu\alpha(\tau)) d\tau; \quad (26)$$

Evolution of density matrix in paths integral formulation

We have

$$\begin{aligned} \rho(\alpha_f^*, \bar{\theta}_f, \alpha'_f, \theta'_f; t_f) = & \int \frac{d^2\alpha'_{in}}{\pi} \frac{d^2\theta'_{in}}{\pi} \frac{d^2\alpha_{in}}{\pi} \frac{d^2\theta_{in}}{\pi} \rho(\alpha_{in}^*, \bar{\theta}_{in}, \alpha'_{in}, \theta'_{in}; t_{in}) \times \\ & \times \mathfrak{D}\alpha^*(\tau) \mathfrak{D}\alpha(\tau) \mathfrak{D}\bar{\theta}(\tau) \mathfrak{D}\theta(\tau) \mathfrak{D}\alpha'^*(\tau) \mathfrak{D}\alpha'(\tau) \mathfrak{D}\bar{\theta}'(\tau) \mathfrak{D}\theta'(\tau) \times \\ & \times \exp \left\{ i \left(S_{full} [\alpha^*(\tau), \alpha(\tau), \bar{\theta}(\tau), \theta(\tau)] - S_{full} [\alpha'^*(\tau), \alpha'(\tau), \bar{\theta}'(\tau), \theta'(\tau)] \right) \right\}, \quad (27) \end{aligned}$$

Fermionic density matrix and influence functional

$$\rho(\bar{\theta}_f, \theta'_f; t_f) = Sp_{\alpha_f=\alpha'_f} \rho(\alpha_f^*, \bar{\theta}_f, \theta'_f, \alpha'_f; t_f) = \int d\theta'_f d\theta_f \mathfrak{D}\bar{\theta}(\tau) \mathfrak{D}\theta(\tau) \mathfrak{D}\bar{\theta}'(\tau) \mathfrak{D}\theta'(\tau) d\theta'_{in} d\theta_{in} \times \\ \times \exp \left\{ i \left(S_f[\bar{\theta}(\tau), \theta(\tau)] - S_f[\bar{\theta}'(\tau), \theta'(\tau)] \right) \right\} F[\theta(\tau), \theta'(\tau)] \quad (28)$$

where $F[\bar{\theta}(\tau), \theta'(\tau)]$ is influence functional of electromagnetic field on fermionic subsystems.

$$F[\bar{\theta}(\tau), \theta'(\tau)] = Sp_{\alpha_f=\alpha'_f} \int \mathfrak{D}\alpha^*(\tau) \mathfrak{D}\alpha(\tau) \mathfrak{D}\alpha'^*(\tau) \mathfrak{D}\alpha'(\tau) \frac{d^2\alpha_{in}}{\pi} \frac{d^2\alpha'_{in}}{\pi} \rho(\alpha_{in}^*, \bar{\theta}_{in}, \alpha'_{in}, \theta'_{in}; t_{in}) \times \\ \times \exp \left\{ i \left(S_b [\alpha^*(\tau), \alpha(\tau)] + S_{int} [\alpha^*(\tau), \alpha(\tau), \bar{\theta}(\tau), \theta(\tau)] - S_b [\alpha'^*(\tau), \alpha'(\tau)] - S_{int} [\alpha'^*(\tau), \alpha'(\tau), \bar{\theta}'(\tau), \theta'(\tau)] \right) \right\} \quad (29)$$

In many cases, we can choose at initial moment t_{in}

$$\rho(\alpha_{in}^*, \bar{\theta}_{in}, \alpha'_{in}, \theta'_{in}; t_{in}) = \rho_f(\bar{\theta}_{in}, \theta'_{in}; t_{in}) \times \rho_b(\alpha_{in}^*, \alpha'_{in}; t_{in}) \quad (30)$$

Influence functional of electromagnetic field

$$F[\bar{\theta}(\tau), \theta'(\tau)] = \int Sp_{\alpha_f = \alpha'_f} \frac{d^2\alpha_{in}}{\pi} \frac{d^2\alpha'_{in}}{\pi} \times \\ \times U_{infl}(\alpha_f^*, \bar{\theta}_f, t_f | \alpha_{in}, \theta_{in}, t_{in}) \rho(\alpha_{in}^*, \bar{\theta}_{in}, \alpha'_{in}, \theta'_{in}; t_{in}) U_{infl}^*(\alpha'_f, \theta'_f, t_f | \alpha'^*_{in}, \bar{\theta}'_{in}; t_{in}) \quad (31)$$

where $U_{infl}(\alpha_f^*, \bar{\theta}_f, t_f | \alpha_{in}, \theta_{in}, t_{in})$ is electromagnetic field transition amplitude from initial state $|\alpha_{in}\rangle$ to final state $|\alpha_f^*\rangle$ inducing by external source j :

$$U_{infl}(\alpha_f^*, \bar{\theta}_f, t_f | \alpha_{in}, \theta_{in}, t_{in}) = \int \mathfrak{D}\alpha^*(\tau) \mathfrak{D}\alpha(\tau) \exp \{iS_{infl}[\alpha^*(\tau), \alpha(\tau), x(\tau)]\} \quad (32)$$

where $S_{infl} [\alpha^*(\tau), \alpha(\tau), \bar{\theta}(\tau), \theta(\tau)] = S_b [\alpha^*(\tau), \alpha(\tau)] + S_{int} [\alpha^*(\tau), \alpha(\tau), \bar{\theta}(\tau), \theta(\tau)]$. In general, influence functional (26) describes the influence (action) of electromagnetic field on fermionic field.

Functional integration over electromagnetic field paths

$$U_{infl}(\alpha_f^*, \bar{\theta}_f, t_f | \alpha_{in}, \theta_{in}, t_{in}) = \exp \left\{ e^{-i\omega(t_f - t_{in})} \alpha_f^* \alpha_{in} - e^2 \int_{t_{in}}^{t_f} \int_{t_{in}}^{\tau} \varepsilon^\mu j_\mu^+(\tau) \varepsilon^{*\nu} j_\nu^-(\tau') e^{i\omega(\tau - \tau')} d\tau d\tau' - \right.$$

$$\left. - i\alpha_{in} e \int_{t_{in}}^{t_f} \varepsilon^\mu j_\mu^+(\tau) e^{-i\omega(\tau - t_{in})} d\tau - i\alpha_f^* e \int_{t_{in}}^{t_f} \varepsilon^{*\mu} j_\mu^-(\tau) e^{-i\omega(t_f - \tau)} d\tau \right\} \quad (33)$$

For multimode field without interaction between modes

$$U_{infl} = \prod_{\mathbf{k}} U_{infl}^{(\mathbf{k})} \quad (34)$$

$$\begin{aligned}
& U_{infl}(\alpha_f^*, \bar{\theta}_f, t_f | \alpha_{in}, \theta_{in}, t_{in}) = \\
&= \prod_{\mathbf{k}, \lambda=1,2} \exp \left\{ e^{-i\omega_{\mathbf{k}}(t_f - t_{in})} \alpha_{\mathbf{k}\lambda}^{(f)*} \alpha_{\mathbf{k}\lambda}^{(in)} - \frac{e^2}{2\omega_{\mathbf{k}} V} \int_{t_{in}}^{t_f} \int_{t_{in}}^{\tau} \varepsilon_{\lambda}^{\mu} j_{\mu}^{+}(\mathbf{k}, \tau) \varepsilon_{\lambda}^{*\nu} j_{\nu}^{-}(\mathbf{k}, \tau') e^{i\omega_{\mathbf{k}}(\tau - \tau')} d\tau d\tau' - \right. \\
&\quad \left. - i\alpha_{\mathbf{k}\lambda}^{(in)} \frac{e}{\sqrt{2\omega_{\mathbf{k}} V}} \int_{t_{in}}^{t_f} \varepsilon_{\lambda}^{\mu} j_{\mu}^{+}(\mathbf{k}, \tau) e^{-i\omega_{\mathbf{k}}(\tau - t_{in})} d\tau - i\alpha_{\mathbf{k}\lambda}^{(f)*} \frac{e}{\sqrt{2\omega_{\mathbf{k}} V}} \int_{t_{in}}^{t_f} \varepsilon_{\lambda}^{*\mu} j_{\mu}^{-}(\mathbf{k}, \tau) e^{-i\omega_{\mathbf{k}}(t_f - \tau)} d\tau \right\} = \\
&= \exp \left\{ \sum_{\mathbf{k}, \lambda} \left(e^{-i\omega_{\mathbf{k}}(t_f - t_{in})} \alpha_f^* \alpha_{in} - \frac{e^2}{2\omega_{\mathbf{k}} V} \int_{t_{in}}^{t_f} \int_{t_{in}}^{\tau} \varepsilon_{\lambda}^{\mu} j_{\mu}^{+}(\mathbf{k}, \tau) \varepsilon_{\lambda}^{*\nu} j_{\nu}^{-}(\mathbf{k}, \tau') e^{i\omega_{\mathbf{k}}(\tau - \tau')} d\tau d\tau' - \right. \right. \\
&\quad \left. \left. - i\alpha_{in} \frac{e}{\sqrt{2\omega_{\mathbf{k}} V}} \int_{t_{in}}^{t_f} \varepsilon_{\lambda}^{\mu} j_{\mu}^{+}(\mathbf{k}, \tau) e^{-i\omega_{\mathbf{k}}(\tau - t_{in})} d\tau - i\alpha_f^* \frac{e}{\sqrt{2\omega_{\mathbf{k}} V}} \int_{t_{in}}^{t_f} \varepsilon_{\lambda}^{*\mu} j_{\mu}^{-}(\mathbf{k}, \tau) e^{-i\omega_{\mathbf{k}}(t_f - \tau)} d\tau \right) \right\} \quad (35)
\end{aligned}$$

Vacuum influence functional

For the case when initial and final states of electromagnetic field are vacuum:

$$\phi_{in}(\alpha_{in}) = \langle \alpha_{in} | 0 \rangle = \exp \left\{ -\frac{1}{2} |\alpha_{in}|^2 \right\}, \quad \phi_f^*(\alpha_f) = \langle 0 | \alpha_f \rangle = \exp \left\{ -\frac{1}{2} |\alpha_f|^2 \right\}. \quad (36)$$

We define influence functional of electromagnetic vacuum

$$F_{(vac|vac)}[\bar{\theta}(\tau), \theta'(\tau)] = \int \frac{d^2 \alpha_f}{\pi} \frac{d^2 \alpha'_f}{\pi} \frac{d^2 \alpha_{in}}{\pi} \frac{d^2 \alpha'_{in}}{\pi} \rho_f(\bar{\theta}_{in}, \theta'_{in}; t_{in}) \times \\ \times \phi_f^*(\alpha_f) U_{infl}(\alpha_f^*, \bar{\theta}_f, t_f | \alpha_{in}, \theta_{in}, t_{in}) \phi_{in}(\alpha_{in}) \phi_{in}^*(\alpha'_{in}) U_{infl}^*(\alpha'_f, \theta'_f, t_f | \alpha'^*_{in}, \bar{\theta}'_{in}; t_{in}) \phi_f^*(\alpha'_f) = \\ = \exp \left\{ - \sum_{\mathbf{k}, \lambda} \frac{e^2}{2\omega_{\mathbf{k}} V} \int_{t_{in}}^{t_f} \int_{t_{in}}^{\tau} \left(\varepsilon_{\lambda}^{\mu} j_{\mu}^+(\mathbf{k}, \tau) \varepsilon_{\lambda}^{*\nu} j_{\nu}^-(\mathbf{k}, \tau') e^{i\omega_{\mathbf{k}}(\tau-\tau')} d\tau d\tau' + \varepsilon_{\lambda}^{\mu} j_{\mu}'^+(\mathbf{k}, \tau) \varepsilon_{\lambda}^{*\nu} j_{\nu}'^-(\mathbf{k}, \tau') e^{-i\omega_{\mathbf{k}}(\tau-\tau')} d\tau d\tau' \right) \right\} \quad (37)$$

From sum over \mathbf{k} to integral: $\sum_{\mathbf{k}} \rightarrow \frac{V}{(2\pi)^3} \int d\mathbf{k}$

$$\begin{aligned}
 & - \sum_{\mathbf{k}} \frac{e^2}{2\omega_{\mathbf{k}} V} \int_{t_{in}}^{t_f} \int_{t_{in}}^{\tau} \left[\sum_{\lambda} \varepsilon_{\lambda}^{\mu} \varepsilon_{\lambda}^{nu} \right] j_{\mu}^{+}(\mathbf{k}, \tau) j_{\nu}^{-}(\mathbf{k}, \tau') e^{i\omega_{\mathbf{k}}(\tau-\tau')} d\tau d\tau' = \\
 & = - \frac{e^2}{(2\pi)^3} \int_{t_{in}}^{t_f} \int_{t_{in}}^{\tau} \int \frac{1}{2\omega_{\mathbf{k}}} \left[\sum_{\lambda} \varepsilon_{\lambda}^{\mu} \varepsilon_{\lambda}^{*\nu} \right] j_{\mu}^{+}(\mathbf{k}, \tau) j_{\nu}^{-}(\mathbf{k}, \tau') e^{i\omega(\tau-\tau')} d\mathbf{x} d\mathbf{x}' d\mathbf{k} d\tau d\tau' = \\
 & = - \frac{e^2}{(2\pi)^3} \int_{t_{in}}^{t_f} \int_{t_{in}}^{\tau} \int \frac{1}{2\omega_{\mathbf{k}}} \left[\sum_{\lambda} \varepsilon_{\lambda}^{\mu} \varepsilon_{\lambda}^{*\nu} \right] j_{\mu}(\mathbf{x}, \tau) j_{\nu}(\mathbf{x}', \tau') e^{-i\mathbf{k}(\mathbf{x}-\mathbf{x}')} e^{i\omega(\tau-\tau')} d\mathbf{x} d\mathbf{x}' d\mathbf{k} d\tau d\tau' = \\
 & = - \frac{e^2}{4\pi i} \int_{t_{in}}^{t_f} \int_{t_{in}}^{\tau} \int \underbrace{\left[\frac{1}{(2\pi)^3} \int \frac{2\pi i d\mathbf{k}}{\omega_{\mathbf{k}}} \left(\sum_{\lambda} \varepsilon_{\lambda}^{\mu} \varepsilon_{\lambda}^{*\nu} \right) e^{-i\mathbf{k}(\mathbf{x}-\mathbf{x}')} e^{i\omega(\tau-\tau')} d\mathbf{k} \right]}_{D^{\mu\nu}(\mathbf{x}-\mathbf{x}', \tau-\tau')} j_{\mu}(\mathbf{x}, \tau) j_{\nu}(\mathbf{x}', \tau') d\mathbf{x} d\mathbf{x}' d\tau d\tau'
 \end{aligned}$$

where $D^{\mu\nu}(\mathbf{x} - \mathbf{x}', \tau - \tau')$ is photon propagator ¹.

$$F_{\langle vac|vac \rangle}[\bar{\theta}(\tau), \theta'(\tau)] = \exp \left\{ -\frac{e^2}{4\pi i} \int_{t_{in}}^{t_f} \int_{t_{in}}^{\tau} \int j_{\mu}(\mathbf{x}, \tau) D^{\mu\nu}(\mathbf{x} - \mathbf{x}', \tau - \tau') j_{\nu}(\mathbf{x}', \tau') d\mathbf{x} d\mathbf{x}' d\tau d\tau' - \right. \\ \left. - \frac{e^2}{4\pi i} \int_{t_{in}}^{t_f} \int_{t_{in}}^{\tau} \int j'_{\mu}(\mathbf{x}, \tau) D^{\mu\nu}(\mathbf{x} - \mathbf{x}', \tau - \tau') j'_{\nu}(\mathbf{x}', \tau') d\mathbf{x} d\mathbf{x}' d\tau d\tau' \right\} \quad (38)$$

For $t_f \rightarrow \infty$, $t_{in} \rightarrow -\infty$ we have relativistic invariant influence functional of electromagnetic vacuum:

$$F_{\langle vac|vac \rangle}[\bar{\theta}(\tau), \theta'(\tau)] = \exp \left\{ -\frac{e^2}{4\pi i} \int \int (j_{\mu}(x) D^{\mu\nu}(x - x') j_{\nu}(x') + j'_{\mu}(x) D^{\mu\nu}(x - x') j'_{\nu}(x')) d^4x d^4x' \right\} \quad (39)$$

So we have effective non-local Lagrangian

$$\mathcal{L} = \bar{\psi}(x)(i\gamma^\mu \partial_\mu - m)\psi(x) + \frac{e^2}{4\pi} j_\mu(x) \int D^{\mu\nu}(x - x') j_\nu(x') dx' \quad (40)$$

Thanks for your attention!