



Recent measurements of K_{13}^{\pm} form factors at NA48

Sergey Shkarovskiy

Joint Institute for Nuclear Research, Dubna

on behalf of the NA48/2 collaboration

QFTHEP'2017


26 June - 3 July, 2017

Yaroslavl, Russia

Outline


- NA48/2 experiment
- K_{13} form factors precision measurement
- Conclusion

NA48/2 kaon beam



2003+2004 ~ 6 months,
 ~ $2 \cdot 10^{11}$ K decays
 Flux ratio: $K^+/K^- \approx 1.8$


Simultaneous K^+ and K^- beams:
 large **charge symmetrization** of
 experimental conditions



NA48/2 detector

Main detector components:

- Magnetic spectrometer (4 DCHs):
4 views/DCH inside a He tank
 $\Delta p/p = 1.02\% \oplus 0.044\% * p$
[p in GeV/c].
- Hodoscope
fast trigger;
precise time measurement (150ps).
- Liquid Krypton EM calorimeter (LKr)
High granularity, quasi-homogenous
 $\sigma_E/E = 3.2\%/E^{1/2} \oplus 9\%/E \oplus 0.42\%$
 $\sigma_x = \sigma_y = 0.42/E^{1/2} \oplus 0.06\text{cm}$
[E in GeV]. (0.15cm@10GeV).
- Hadron calorimeter, muon veto counters, photon vetoes.



$K^\pm \rightarrow \pi^0 l^\pm \nu$ (K_{l3}^\pm) form factors

exper. input for $|V_{us}|$ extraction (apart from $\Gamma(K_{l3}^\pm)$)

Without radiative effects : $\rho_0 = d^2 N / (dE_l dE_\pi) \sim A f_+^2(t) + B f_+(t) f_-(t) + C f_-^2(t)$, where

$$t = (P_K - P_\pi)^2 = M_K^2 + M_\pi^2 - 2 M_K E_\pi$$

$f_-(t) = (f_+(t) - f_0(t))(m_K^2 - m_\pi^2)/t$. (just another formulation, f_0 is «scalar» and f_+ is «vector» FF),

E_l is charged lepton energy, E_π is π^0 energy (both in the kaon rest frame).

$$A = M_K(2 E_l E_\nu - M_K(E_\pi^{\max} - E_\pi)) + M_l^2 ((E_\pi^{\max} - E_\pi)/4 - E_\nu)$$

$$B = M_l^2 (E_\nu - (E_\pi^{\max} - E_\pi)/2) \quad \text{negligible for Ke3}$$

$$C = M_l^2 (E_\pi^{\max} - E_\pi)/4 \quad \text{negligible for Ke3}$$

$$E_\pi^{\max} = (M_K^2 + M_\pi^2 - M_l^2)/(2 M_K)$$

FF Parameterisation (PDG name)	$f_+(t, \text{parameters})$	$f_0(t, \text{parameters})$
Quadratic (linear for $\bar{f}_0(t)$)	$1 + \lambda'_+ t/m_\pi^2 + 1/2 \lambda''_+ (t/m_\pi^2)^2$	$1 + \lambda'_0 t/m_\pi^2$
Pole	$M_V^2 / (M_V^2 - t)$	$M_S^2 / (M_S^2 - t)$
Dispersive* H(t), G(t): functions fixed from theory and other experiments. Depend on 2 (H) and 3 (G) extra external parameters known with a given* uncertainty.	$\exp((\Lambda_+ + H(t)) t/m_\pi^2)$	$\exp((\ln[C] - G(t)) t/(m_K^2 - m_\pi^2))$

* [V. Bernard, M. Oertel, E. Passemar, J. Stern. Phys.Rev. D80 (2009) 034034]

We use MC **radiative** decay generator of C.Gatti [Eur.Phys.J. C45 (2006) 417–420] provided by KLOE collaboration. It includes $f_0 = f_+ = 1 + \lambda'_+ t/m_\pi^2$.

Data: 16 special runs from the NA48/2 data taken in 2004 (3 days)

Trigger: 1 charged track (2 hodoscope hits) and $E_{LKr} > 10$ GeV

Registered :

- **1 track** (> 0 candidates): $P_e \geq 5$ GeV, $P_\mu \geq 10$ GeV , $R_{MUV} > 30$ cm, $|X_{MUV}, Y_{MUV}| < 115$ cm.
- **2 LKr clusters** (> 1 candidates): $E > 3$ GeV, to closest track > 15 cm.

Neutrino is missing, beam geometry and average momentum P_b are measured from $K^\pm \rightarrow \pi^\pm \pi^+ \pi^-$

Kaon momentum reconstruction


Two solutions of the quadratic equation for P_K :

$$P_{1,2} = (\phi P_{Zb} \pm \text{SQRT}(d)) / (E^2 - P_{Zb}^2), \text{ where}$$

$$\phi = 0.5 (M_K^2 + E^2 - P_t^2 - P_{Zb}^2),$$

$$d = (\phi^2 P_{Zb}^2 - (E^2 - P_{Zb}^2)(M_K^2 E^2 - \phi^2))$$

When $d < 0$, we assume $d=0$.



- Best P_K solution = closest $P_{1,2}$ to the average beam momentum P_b measured from $3\pi^\pm$ decays for each run is used to choose the.
- A cut: -7.5 GeV/c $< (P_K - P_b) < 7.5$ GeV/c
- For each event, separately for K_{e3} and $K_{\mu3}$ selections, the combination with a minimum $\Delta P = |P_K - P_b|$ is the best candidate.

Selection


π^0 :

- A pair of clusters in-time (within **5 ns**) without any in-time extra clusters
- Distance between the clusters in a pair **> 20 cm**
- **$E(\pi^0) > 15 \text{ GeV}$** (for the trigger efficiency)
- Z of decay: from 2γ assuming π^0 mass («neutral Z»)
- DCH1 inner flange cut for the both γ

Track selection and identification

- A good track in-time with the π^0 within 10 ns.
- No extra good track within 8 ns (against showers).
- If **$2.0 > E_{\text{LKr}}/P_{\text{DCH}} > 0.9$** , it is **an electron of K_{e3}** .
- If **$E_{\text{LKr}}/P_{\text{DCH}} < 0.9$** (for true muons it cuts nothing) and there is a MUV muon associated, it is **a $K_{\mu 3}$ muon**.

Decay vertex reconstruction



Reminder: Preliminary result reported in 2012 was based on the «charged» vertex definition (from CDA between the track and the beam), that leads to high sensitivity to the exact beam shape simulation (due to the systematic shift of the vertex closer to beam).

Neutral vertex is chosen finally

(no transverse bias): $Z_{\text{decay}} = Z(\pi^0)$;

$X_{\text{decay}}, Y_{\text{decay}} =$ impact point of reconstructed charged track on Z_{decay} plane

Final cuts

For K_{e3}

- ν transversal momentum with respect to beam axis $P_t \geq 0.03$ GeV against $K^\pm \rightarrow \pi^\pm \pi^0$ with π^\pm misidentified as e (when $E/P > 0.9$);
- $P_L(\nu)^2 = (E\nu)^2/c^2 - (P_t\nu)^2 > 0.0014$ GeV²/c²
[negative tail and zero region sensitive to beam shape]

For $K_{\mu 3}$

- against the background from $K^\pm \rightarrow \pi^\pm \pi^0$ with $\pi^\pm \rightarrow \mu^\pm \bar{\nu}$
 $m(\pi^+ \pi^0) < 0.47$ GeV/c²
 $m(\pi^+ \pi^0) < (0.6 - P_t(\pi^0))$ GeV/c²
 $m(\mu^\pm \bar{\nu}) > 0.18$ GeV/c² (to exclude π^+ mass region)
- a cut against $\pi^\pm \pi^0 \pi^0$: $(P_2 - P_1) < 60$ GeV
[a difference between two P solution is large when one π^0 is missing]

For both K_{l3}

Beam transverse elliptic variable $B_{ell} < 11$.

X_n, Y_n, Z_n are the reconstructed neutral vertex coordinates, X_n^0, Y_n^0 , $\sigma_{X_n}, \sigma_{Y_n}$ are the reconstructed beam central positions and widths with respect to the run-dependent beam axis Z_b .

Background

Decay	Notation	Br, %	Ng, 10^6	$F_e, 10^{-3}$	$F_\mu, 10^{-3}$
$K^\pm \rightarrow \pi^\pm (\pi^0 \rightarrow 2\gamma)$	2π	20.66	393.2	0.270	0.264
$K^\pm \rightarrow \pi^\pm 2(\pi^0 \rightarrow 2\gamma)$	3π	1.761	62.5	0.286	1.833
$K^\pm \rightarrow \pi^\pm (\pi^0 \rightarrow e^+e^-\gamma)$	$2\pi D$	1.174	1.5	0.049	0.000
$K^\pm \rightarrow \pi^\pm \gamma (\pi^0 \rightarrow 2\gamma)$	$2\pi\gamma$	0.0275	35.3	0.004	0.044
$K^\pm \rightarrow \pi^0 \mu^\pm \nu (\mu \rightarrow e\nu)$	$K_{\mu 3}^e$	0.03353	174.3	0.004	0.000

Br — branching ration

Ng — number of MC generated events

F_e — estimated background contamination in K_{e3} data

F_μ — estimated background contamination in $K_{\mu 3}$ data

Events-weighting fit procedure

- Experimental Dalitz plot is corrected for the simulated background.
- For each fit iteration, the model Dalitz plot is filled in with an MC simulated reconstructed center-of-mass pion and lepton energies. Each event is weighted by

$$w = \rho_0(E_\pi^{\text{true}}, E_l^{\text{true}}, FF_{\text{fit}}) / \rho_0(E_\pi^{\text{true}}, E_l^{\text{true}}, FF_{\text{MC generator}}),$$

where ρ_0 is the non-radiative Dalitz density formula.





- MINUIT package is searching for the FF_{fit} parameters minimizing the standard χ^2 value:

$$\chi^2 = \sum_{i,j} \frac{(D_{i,j} - MC_{i,j})^2}{(\delta D_{i,j})^2 + (\delta MC_{i,j})^2},$$

where i,j means the Dalitz plot cell indices, $D_{i,j}$ is the background-corrected experimental data content of the cell, $MC_{i,j}$ is the weighted MC bin content, and $\delta D_{i,j}$, $\delta MC_{i,j}$ are the corresponding statistical errors.



At least 20 data events per cell are required in the fit area, so χ^2 works well.

Experimental Dalitz plots and fits areas (5x5 MeV cells)





Dalitz plot projections


- Data-Bg
- + MC fit result (quadr.)
- + (Data-Bg)/MC






K_{e3}



Marginally significant slope within the radiative correction precision. Radiative effect uncertainty is taken into account as a contribution to systematics.



$K_{\mu 3}$



Small deviation in the Bg area. Bg-related uncertainty is included into syst. error.

Results for the joint K_{l3} analysis

Analysis has been performed:

- For K_{e3}
- For $K_{\mu3}$
- For the combined K_{l3} result

Quadratic parameterization
(in units of 10^{-3})

$$\chi^2/\text{ndf} = 1004.6/1073$$

Correlation coefficients

	$\lambda''_+(K_{l3})$	$\lambda_0(K_{l3})$
$\lambda'_+(K_{l3})$	-0.954	-0.076
$\lambda''_+(K_{l3})$		0.035

	$\lambda'_+(K_{l3})$	$\lambda''_+(K_{l3})$	$\lambda_0(K_{l3})$
Central values	23.35	1.73	14.90
Stat. error	0.75	0.29	0.55
Beam scattering	0.90	0.35	0.45
LKr nonlinearity	0.19	0.03	0.35
LKr scale	0.66	0.15	0.08
Background	0.07	0.03	0.04
Trigger	0.20	0.10	0.45
Accidentals	0.23	0.08	0.08
Acceptance	0.24	0.07	0.01
Pk average	0.04	0.01	0.24
Pk spectra	0.01	0.00	0.04
Neutrino P cut	0.18	0.04	0.03
Binning	0.08	0.02	0.16
Resolution	0.00	0.02	0.14
Radiative	0.22	0.01	0.06
Syst. error	1.23	0.41	0.80
Total error	1.44	0.50	0.97

Pole parameterization (in MeV)

$$\chi^2/\text{ndf} = 1001.1/1074$$

	$m_V(K_{l3})$	$m_S(K_{l3})$
Central values	894.3	1185.5
Stat. error	3.2	16.6
Beam scattering	0.1	27.2
LKr nonlinearity	1.7	14.3
LKr scale	3.9	3.6
Background	0.1	0.6
Trigger	0.7	12.9
Accidentals	0.5	0.0
Acceptance	0.7	3.3
Pk average	0.3	8.8
Pk spectra	0.1	1.5
Neutrino P cut	1.0	0.7
Binning	0.4	5.1
Resolution	0.8	4.3
Radiative	2.7	2.7
Syst. error	5.4	35.5
Total error	6.3	39.2

Correlation = - 0.278

Dispersive parameterization (in units of 10^{-3})




$$\chi^2/\text{ndf} = 998.3/1074 \text{ (the best)}$$

	$\Lambda_+(K_{l3})$	$\ln[C](K_{l3})$
Central values	22.67	189.12
Stat. error	0.18	4.91
Beam scattering	0.01	8.39
LKr nonlinearity	0.10	4.04
LKr scale	0.23	0.88
Background	0.00	0.14
Trigger	0.04	3.73
Accidentals	0.03	0.01
Acceptance	0.04	0.92
Pk average	0.02	2.63
Pk spectra	0.00	0.44
Neutrino P cut	0.06	0.16
Binning	0.03	1.46
Resolution	0.05	1.28
Radiative	0.16	0.75
Parameterization	0.44	3.04
Syst. error	0.55	11.09
Total error	0.58	12.13

Correlation = - 0.035

Joint K_{13} results

- Comparison for quadratic fit: λ'_+ , λ''_+ , λ'_0
- Parameter correlation 1σ ellipses rather than 68% for better visibility



Conclusion

- K_{l3} form factors measurement is performed by NA48/2 on the basis of 2004 run selected $4.28 \cdot 10^6$ (K_{e3}) and $2.91 \cdot 10^6$ ($K_{\mu3}$) events.
- Result is competitive with the other ones in $K_{\mu3}$ mode, and a smallest error in K_{e3} has been reached, that gives us also the most precise combined K_{l3} result.
- For the first time both K^+ and K^- K_{e3} decays were studied together.

Spares







B_{ell} distributions in a wide area

~ 3σ range is relatively well simulated as well as the very far tail.

But the discrepancy near ~5-10 is not described by the known background.


Sensitivity to the background variation at the very far tail (>20) is used to measure the Bg-related systematic uncertainty.

It looks like a small wide component of the beam, that becomes negligible for $B_{ell} > 11$. For wider cuts final results are stable.




B_{ell} distributions with the modified MC beam (systematics)

Data $3\pi^\pm$ decay: Kaon impact points X,Y at the focus plane



Focused scattering simulated in MC: 3% of beam kaons are additionally scattered into a series of rings with a different radius at focus > 2.2 cm.



This MC simplified modification is not used for the FF central values extraction (only for systematics estimate). So we need a wide radius cut to avoid the acceptance distortion, and also we need a vertex reconstruction, that is not too sensitive to the transverse general shift of the decay — it is a Neutral vertex rather than CDA.

Selection

Min bias **trigger**: 1 track and $E_{\text{LKr}} > 10 \text{ GeV}$ ((sevt->trigWord >> 11) & 1)

N of good clusters > 1 :

- LKr standard nonlinearity correction for Data clusters (user_lkr_calcor_SC)
- LKr small final nonlinearity correction for MC clusters, extracted from $\pi^+\pi^0\pi^0$ (see April 2007 talk of Di Lella and Madigozhin)
- LKr scale corrections from K_{e3} E/P (different for Data and MC, sub-permill precision)
- Cluster status ≤ 4
- Cluster energy $\geq 3 \text{ GeV}$
- Distance to dead cell $\geq 2 \text{ cm}$
- Radius at LKr $\geq 15 \text{ cm}$
- In standard LKr acceptance
- Distance to any in-time (within 10 ns) track impact point at LKr $\geq 15 \text{ cm}$
- Distance to any another in-time (within 5 ns) cluster $\geq 10 \text{ cm}$

In Monte Carlo everything is in-time

N of good tracks > 0 :

- $P_e \geq 5 \text{ GeV}$, $P_\mu \geq 10 \text{ GeV}$ (muon case cut applied after identification)
- Track momenta α, β corrections both for data and MC
- If there is the associated LKr cluster, its cluster status ≤ 4
- Track quality ≥ 0.6
- Distance to dead cell $\geq 2 \text{ cm}$
- Radius at every DCH(1,2,3,4) $\geq 15 \text{ cm}$
- **Reject DCH tracks with $0 \text{ cm} < X(\text{DCH4}) < 6 \text{ cm}$ && $Y(\text{DCH4}) > 0$** (inefficient band)
- **$K_{\mu 3}$ DCH track: for all 3 MUV planes $R_{\text{MUV}} > 30 \text{ cm}$, $|X_{\text{MUV}}, Y_{\text{MUV}}| < 115 \text{ cm}$.**
- LKr impact point is in LKr acceptance

π^0 selection

- Check all the pairs of good in-time (within 5 ns) clusters
- Calculate π^0 time t_π (average of two γ ones) and reject the combination, if there is a good extra cluster in 5 nanoseconds around t_π (to suppress $\pi^+\pi^0\pi^0$ and showers).
- Make the projectivity correction for the experimental data and MC.
- Reject the pair, if the distance between the clusters is < 20 cm
- $E_{\pi^0} > 15$ GeV (for trigger efficiency: trigger E LKr > 10 GeV).
- Calculate Z_n from two γ , assuming π^0 mass
- -1600 cm $< Z < 9000$ cm
- DCH flunge gamma cut for the both γ

Track selection and identification

For each found good π^0 check all the good tracks:

- In-time with π^0 (within 10 ns)
- There is no extra good track within 8 ns around the track time (against showers).
- If $2.0 > E/P > 0.9$, it is an electron (K_{e3})
- If $E/P < 0.9$ and there is a muon associated, it is a muon ($K_{\mu3}$)

First iteration decay vertex position:

- $Z_{\text{decay}} = Z(\pi^0)$
- $X_{\text{decay}}, Y_{\text{decay}} =$ impact point of reconstructed charged track on the transversal plane, defined by Z_{decay}

Blue field correction:

With the «first iteration vertex», we implement the Blue field correction, obtain corrected track slopes and recalculate vertex X,Y again.

Beam position correction:

We know the position of beam axis in space (it is always displaced slightly from the nominal Z axis). For the CMC tuning, these positions were measured for each run from $3\pi^\pm$ data many years ago.

We use these data to calculate all the relevant values with respect to the current run beam axis rather than with respect to nominal Z arrow. First of all, we calculate the vertex (x,y) with respect to the beam center X_b, Y_b at this Z_n .

Vertex position cut (very wide):

$$\text{SQRT}(((X-a_X(Z))/\sigma_X(Z))^2 + ((Y-a_Y(Z))/\sigma_Y(Z))^2) < \mathbf{11.0}$$

Here a_X, a_Y, σ_X and σ_Y are the functions of Z and represent the average position and width of the beam with respect to standard ($3\pi^\pm$) beam position.

They are obtained by Gaussian fit (± 1.2 cm around maximum) for Z slices, separately for MC and Data, for X and Y and for positive and negative beams. Then these points are parametrised as functions of Z by polinomes of 5-th degree of Z.

Final stage of the selection

- $P_{\perp}(\nu)^2 > 0.0014 \text{ GeV}^2$ for K_{e3} only
- Quadratic equation for P_K is solved, if no solutions, the combination is taken with zero discriminant. With the above $P_{\perp}(\nu)^2$ requirement, such a cases are rare for K_{e3} .
- Average beam momentum P_b measured from $3\pi^{\pm}$ decays for each run is used to choose the best P_K solution (closest to P_b from two ones).
- $-7.5 \text{ GeV}/c < (P_K - P_b) < 7.5 \text{ GeV}/c$



- For $K_{\mu3}$, the cut against $K^{\pm} \rightarrow \pi^{\pm}\pi^0$ with $\pi^{\pm} \rightarrow \mu^{\pm}\bar{\nu}$:
 $m(\pi^+\pi^0) < 0.47 \text{ GeV}$ and $m(\pi^+\pi^0) < (0.6 - P_t(\pi^0)) \text{ GeV}$;
- For $K_{\mu3}$, one more cut against $K^{\pm} \rightarrow \pi^{\pm}\pi^0$ with $\pi^{\pm} \rightarrow \mu^{\pm}\bar{\nu}$:
 $m(\mu^{\pm}\bar{\nu}) > 0.18 \text{ GeV}$;
- For $K_{\mu3}$ only: a cut against $\pi^{\pm}\pi^0\pi^0$: $(P_2 - P_1) < 60 \text{ GeV}$
 \Leftrightarrow in terms of P_K equation discriminant squared $d = ((P_2 - P_1)/2)^2$: $d < 900 \text{ GeV}^2$;
- For K_{e3} , the ν transversal momentum with respect to beam axis must be
 $P_t \geq 0.03 \text{ GeV}$: a cut against $K^{\pm} \rightarrow \pi^{\pm}\pi^0$ with π^{\pm} misidentified as e (when $E/P > 0.9$).

In every event, separately for K_{e3} and $K_{\mu3}$, the combination with the minimum $\Delta P = |P_K - P_b|$ is chosen as the best candidate.

A complex nature of $(P_L^v)^2$ - dependent K_{e3} systematic effect

- 1) Mismeasurement of decay transversal coordinates happens (in the neutral vertex case it also involves the LKr clusters mismeasurement).
- 2) As a consequence, a small mismeasurement of transversal $(P_t^v)^2$
- 3) As a consequence, a small mismeasurement of $(P_L^v)^2 = (E^v)^2 - (P_t^v)^2$
- 4) As a consequence, a small mismeasurement of $D = ((P_1^K - P_2^K)/2)^2$
- 5) When D itself is small or negative, even small D mismeasurement is relatively not small.
- 6) Distorted D changes in a different way the probability of the «best» P^K choice (we take the closest to average true $\langle P^K \rangle$) for different vertex definitions and for MC and Data, depending on true P^K spectrum. The wrong choice may also depend on the correlations between true P^K and the transversal decay coordinates.
- 7) Mistake in P^K choice from two options may be not small, it is of the order of spectrum width (few GeV), and it leads to relatively big mismeasurement of Dalits plot variables, especially for E_π^* .
 - Correct simulation of this effect seems to be difficult, we have only a simple beam correction for the scattered component.
 - But we know, where the problem is concentrated (small $(P_L^v)^2$), so we just cut the problematic region.

For $K_{\mu 3}$ only: a cut against $\pi^{\pm}\pi^0\pi^0$: $(P_2-P_1)<60$ GeV \Leftrightarrow $D = ((P_2-P_1)/2)^2 < 900$ GeV²




Equally normalized distributions of signal and background events are shown in order to check that the cut is doing its work in both cases.


But the absolute K_{e3} background level is much smaller than for $K_{\mu 3}$.
So we don't use this cut for K_{e3} and save some experimental statistics.

For K_{e3} , the ν transversal momentum with respect to beam axis must be $P_t \geq 0.03$ GeV.

It is a cut against $K^\pm \rightarrow \pi^\pm \pi^0$ with π^\pm misidentified as e (when $E/P > 0.9$).



Cuts for $K_{\mu 3}$ against the background from $K^{\pm} \rightarrow \pi^{\pm}\pi^0$ with $\pi^{\pm} \rightarrow \mu^{\pm}\bar{\nu}$



$m(\pi^+\pi^0) < 0.47 \text{ GeV}/c^2$ and
 $m(\pi^+\pi^0) < (0.6 - P_t(\pi^0)) \text{ GeV}/c^2$

$m(\mu^{\pm}\bar{\nu}) > 0.18 \text{ GeV}/c^2$
 (to exclude π^+ mass region)





K_{e3} requirement: $P_L(v)^2 > 0.0014 \text{ GeV}^2$

$(P_L v)^2$ normalized distributions
(Data and MC with background)


$P_L(v)^2 = (E^v)^2 - (P_t^v)^2$
negative tail is difficult to simulate precisely, as it depends on the beam transverse shape (scattering) via P_t^v .

For K_{e3} only the region of small and negative $P_L(v)^2$ induces a systematic FF uncertainty ($P_L(v)^2$ dependence), that is avoided by this cut.

Peak sharpness residual mismatch is used to check $(P_L v)^2$ resolution systematics related to this cut.



Neutral Z normalized distributions comparison



Residual discrepancy ($\sim 1\%$) is taken into account as a contribution to systematic uncertainty = variation of final result due to the change of geometrical acceptance by the factor of 1.002, that corrects the K_{e3} differences.

Experimental systematics

<i>Contribution</i>	<i>Approach to the uncertainty calculation</i>
Beam scattering	Effect of the additional beam fraction imitating the beam scattering
LKr nonlinearity	Effect of the final nonlinearity correction
LKr scale	Effect of the LKr scale shift allowed by Data/MC electron E/P peak
Background	Effect of the background contribution change within B_{ell} distribution tails Data/MC agreement. It absorbs the PDG branching fraction errors
Trigger efficiency	Effect of the measured quadratically smoothed trigger efficiency (~100%)
Accidentals	Effect of the time windows doubling for clusters and tracks acceptance
Acceptance	Effect of small transversal detector cuts increasing for MC, that (over) corrects Z distributions
P_K average	Effect of beam $\langle P_K \rangle$ possible mismeasurement
P_K spectra	Effect of the MC true P_K spectra variation within the agreement of measured MC/Data P_K spectra
Neutrino P cut	Effect of the artificial $(P_L^\nu)^2$ resolution variation within $(P_L^\nu)^2$ peak sharpness MC/Data agreement
Binning	Effect of the bins doubling for the both Dalitz plot dimensions
Resolution	Difference between the main events weighting approach and the acceptance correction technique that is more sensitive to resolution

External contributions to systematic uncertainty.

<i>Contribution</i>	<i>Approach to the uncertainty calculation</i>
Radiative correction precision	Effect of the theoretical uncertainty in the radiative Dalitz plot <i>corrections</i> in terms of one-dimensional slopes.
Parameterization for Dispersive fits	100 fits with the independently sampled 5 external parameters known with a given uncertainty.

The full analysis is performed and form factor parameters are extracted:

- For K_{e3}
- For $K_{\mu3}$
- For the combined K_{l3} result: A joint fits are done by minimizing of the sum $\chi^2(K_{e3}) + \chi^2(K_{\mu3})$ with a common set of fit parameters. This is repeated also for each of the systematic uncertainty studies.

LKr Nonlinearity

Use 2004 $\pi^0\pi^0\pi^{+-}$ data
(done for cusp analysis):

$22 < E(\pi^0_1) < 26$ GeV
 $E(\pi^0_2) < E(\pi^0_1)$
 $E(\gamma)^{\max} < 0.55 E(\pi^0)$ for both π^0

Final correction for MC:



- P_0 1.0170
- P_1 -0.48025E-02
- P_2 0.45538E-03
- P_3 -0.14474E-04

E: cluster energy in GeV

$$f = P_0 + P_1 E + P_2 E^2 + P_3 E^3$$

if $(f > 1)$ $E = E/f$

100% of the final correction effect is taken as the nonlinearity-related uncertainty.



Blue field correction:

With the «first iteration vertex», we implement the Blue field correction, obtain corrected track slopes and recalculate vertex X,Y again.

Beam position correction:

We know the position of beam axis in space (it is always displaced slightly from the nominal Z axis). For the CMC tuning, these positions were measured for each run from $3\pi^\pm$ data many years ago.

We use these data to calculate all the relevant values with respect to the **current run beam axis** rather than with respect to nominal Z arrow. First of all, we calculate the vertex (x,y) with respect to the beam center X_b, Y_b at this Z_n .

Vertex position cut (very wide):

$$\text{SQRT}(((X-a_X(Z))/\sigma_X(Z))^2 + ((Y-a_Y(Z))/\sigma_Y(Z))^2) < \mathbf{11.0}$$

Here a_X, a_Y, σ_X and σ_Y are the functions of Z and represent the average position and width of the beam with respect to standard ($3\pi^\pm$) beam position.

They are obtained by Gaussian fit (± 1.2 cm around maximum) for Z slices, separately for MC and Data, for X and Y and for positive and negative beams. Then these points are parametrised as functions of Z by polinomes of 5-th degree of Z.

Results for K_{e3} and $K_{\mu3}$

$\chi^2/\text{NDF}(K_{e3})$:
609.4/687

$\chi^2/\text{NDF}(K_{\mu3})$:
391.2/384

Quadratic
parameterization
(in units of 10^{-3})

	$\lambda'_+(K_{e3})$	$\lambda''_+(K_{e3})$	$\lambda'_+(K_{\mu3})$	$\lambda''_+(K_{\mu3})$	$\lambda_0(K_{\mu3})$
Central values	23.52	1.60	23.32	2.14	14.33
Stat. error	0.78	0.30	3.08	1.06	1.11
Beam scattering	0.90	0.32	0.25	0.12	0.58
LKr nonlinearity	0.28	0.01	2.85	0.73	0.93
LKr scale	0.68	0.12	0.83	0.18	0.14
Background	0.07	0.04	0.26	0.05	0.04
Trigger	0.27	0.13	0.67	0.23	0.12
Accidentals	0.24	0.08	0.01	0.00	0.01
Acceptance	0.28	0.08	0.85	0.23	0.25
Pk average	0.06	0.01	0.20	0.07	0.32
Pk spectra	0.00	0.00	0.12	0.04	0.00
Neutrino P cut	0.18	0.04	0.00	0.00	0.00
Binning	0.05	0.00	0.11	0.05	0.15
Resolution	0.01	0.02	1.44	0.46	0.39
Radiative	0.20	0.01	0.15	0.03	0.06
Syst. error	1.29	0.39	3.50	0.96	1.25
Total error	1.51	0.49	4.67	1.43	1.67

Correlation
-0.927

Correlation

	$\lambda''_+(K_{\mu3})$	$\lambda_0(K_{\mu3})$
$\lambda'_+(K_{\mu3})$	-0.969	0.851
$\lambda''_+(K_{\mu3})$		-0.810

26 June- 3 July, 2017

QFTHEP'2017, Yaroslavl, Russia

Pole parameterization (in units of 10^{-3})

Dispersion parameterization (in units of 10^{-3})

$\chi^2/\text{NDF}(K_{e3})$:
609.3/688

$\chi^2/\text{NDF}(K_{\mu3})$:
388.0/385

$\chi^2/\text{NDF}(K_{e3})$:
609.1/688

$\chi^2/\text{NDF}(K_{\mu3})$:
385.8/385

	$m_V(K_{e3})$	$m_V(K_{\mu3})$	$m_S(K_{\mu3})$
Central values	896.8	879.1	1196.4
Stat. error	3.4	8.1	18.1
Beam scattering	1.4	7.6	22.6
LKr nonlinearity	3.5	9.6	6.2
LKr scale	5.3	4.1	2.2
Background	0.4	1.5	0.7
Trigger	0.8	0.1	12.7
Accidentals	0.5	0.0	0.3
Acceptance	1.3	2.4	1.0
Pk average	0.3	0.2	9.0
Pk spectra	0.1	0.0	1.6
Neutrino P cut	1.2	0.0	0.0
Binning	0.7	0.5	4.5
Resolution	0.6	2.2	1.0
Radiative	3.2	0.8	1.6
Syst. error	7.6	13.5	28.8
Total error	8.3	15.7	34.0

	$\Lambda_+(K_{e3})$	$\Lambda_+(K_{\mu3})$	$\ln[C](K_{\mu3})$
Central values	22.54	23.55	186.68
Stat. error	0.20	0.50	5.12
Beam scattering	0.09	0.48	7.05
LKr nonlinearity	0.20	0.60	2.08
LKr scale	0.31	0.26	0.50
Background	0.02	0.10	0.15
Trigger	0.04	0.01	3.62
Accidentals	0.03	0.00	0.09
Acceptance	0.08	0.16	0.35
Pk average	0.02	0.01	2.62
Pk spectra	0.00	0.00	0.46
Neutrino P cut	0.07	0.00	0.00
Binning	0.04	0.03	1.24
Resolution	0.03	0.10	0.50
Radiative	0.18	0.05	0.49
Parameterization	0.44	0.49	2.95
Syst. error	0.62	0.97	9.23
Total error	0.65	1.10	10.55

26 June- 3 July, 2017

Correlation
OF THEP'2017, Yaroslavl, Russia
0.320

Correlation
0.408